

## Recommended Immunization Schedule for Persons Aged 7 Through 18 Years—United States • 2010

For those who fall behind or start late, see the schedule below and the catch-up schedule

Vaccine ▼	Age ►	7–10 years	11–12 years	13–18 years
Tetanus, Diphtheria, Pertussis <sup>1</sup>			Tdap	Tdap
Human Papillomavirus <sup>2</sup>	<i>see footnote 2</i>		HPV (3 doses)	HPV series
Meningococcal <sup>3</sup>		MCV	MCV	MCV
Influenza <sup>4</sup>		Influenza (Yearly)		
Pneumococcal <sup>5</sup>		PPSV		
Hepatitis A <sup>6</sup>		HepA Series		
Hepatitis B <sup>7</sup>		Hep B Series		
Inactivated Poliovirus <sup>8</sup>		IPV Series		
Measles, Mumps, Rubella <sup>9</sup>		MMR Series		
Varicella <sup>10</sup>		Varicella Series		

Range of recommended ages for all children except certain high-risk groups

Range of recommended ages for catch-up immunization

Range of recommended ages for certain high-risk groups

This schedule includes recommendations in effect as of December 15, 2009. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Considerations should include provider assessment, patient preference, and the potential for adverse events. Providers should consult the relevant Advisory

Committee on Immunization Practices statement for detailed recommendations: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS) at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.